



Depression and an ageing population



Major depressive disorder (MDD) and dementia commonly co-occur within the ageing population and their overlap is complex^{1,2}





MDD is a risk factor for dementia, although depression can appear years before other symptoms of dementia³




Patients with depressive symptoms have a **1.2–2.4x** increased risk of developing dementia^{4–8}

MDD and co-morbid dementia can reduce QoL and have a negative impact on caregiver burden^{9,10}



Patients with co-morbid depression and dementia have a **54%** lower QoL than the general older adult population⁹

Inadequate treatment of MDD and co-morbid dementia can lead to:^{11–15}



Non-pharmacological approaches in the treatment of depressive symptoms in dementia:^{16,17}



Treating patients with MDD and co-morbid dementia can be challenging¹⁷⁻²⁰



Depression with Alzheimer's disease is under-diagnosed as:¹⁷



Cognitive symptoms related to depression **undermine neuropsychological assessment**¹⁹



MDD **features/functional impairments** may be **less noticeable**¹⁸




Symptoms of depression are **underestimated**²⁰

There is **limited evidence** supporting the efficacy of antidepressants in patients with MDD and co-morbid dementia²¹



Depression **should not** automatically be assumed to be an **inevitable “part of dementia”**²²



References

1. Asmer MS et al. J Clin Psychiatry 2018;79:17r11772
2. Weisenbach SL et al. Curr Psychiatry Rep 2012;14:280–8
3. Byers AL, Yaffe K. Nat Rev Neurol 2011;7:323–31
4. Alzheimer's Disease International. World Alzheimer Report 2014. Available at: <https://www.alzint.org/u/WorldAlzheimerReport2014.pdf>. Accessed 3 October 2023
5. Barnes DE et al. Arch Gen Psychiatry 2012;69:493–8
6. Santabárbara J et al. Actas Esp Psiquiatr 2020;48:169–80
7. Almeida OP et al. Transl Psychiatry 2017;7:e1117
8. Holmquist S et al. PLoS Med 2020;17:e1003016
9. Winter Y et al. J Neurosci Rural Pract 2011;2:27–32
10. Martire LM et al. J Am Geriatr Soc 2010;58:23–9
11. Curran EM, Loi S. MJA Open 2012;1(Suppl 4):40–3

12. Shim YS, Yang DW. Arch Gerontol Geriatr 2006;43:277–83
13. Rapp MA et al. Am J Geriatr Psychiatry 2011;19:357–63
14. Perna L et al. Soc Psychiatry Psychiatr Epidemiol 2019;54:793–801
15. Günak MM et al. JAMA Psychiatry 2021;78:659–66
16. Watt JA et al. BMJ 2021;372:n532
17. Goodarzi Z et al. BMC Neurol 2016;16:244
18. Tetsuka S. Aging Dis 2021;12:1920–34
19. Agüera-Ortiz L et al. Front Psychiatry 2021;12:638651
20. Novais F, Starkstein S. J Alzheimer's Dis 2015;47:845–55
21. Dudas R et al. Cochrane Database Syst Rev 2018;8:CD003944
22. RANZCP. Use of antidepressants to treat depression in dementia. Position statement. 2015. Available at: <https://www.ranzcp.org/news-policy/policy-and-advocacy/position-statements/antidepressants-to-treat-depression-in-dementia>. Accessed 3 Oct 2023